



E1546

JACC March 27, 2012

Volume 59, Issue 13



## Pericardial/Myocardial Disease

### NON-INVASIVE IDENTIFICATION OF SENILE SYSTEMIC AMYLOIDOSIS: INCREMENTAL DIAGNOSTIC ROLE OF 99mTc-DPD SCINTIGRAPHY

ACC Moderated Poster Contributions

McCormick Place South, Hall A

Sunday, March 25, 2012, 9:30 a.m.-10:30 a.m.

Session Title: Pericardial/Myocardial Disease I

Abstract Category: 12. Pericardial/Myocardial Disease

Presentation Number: 1125-32

Authors: *Candida Cristina Quarta, Pier Luigi Guidalotti, Simone Longhi, Cinzia Pettinato, Ornella Leone, Fabrizio Salvi, Alessandra Ferlini, Agnese Milandri, Pamela Gallo, Christian Gagliardi, Elena Biagini, Claudio Rapezzi, University of Bologna, S.Orsola-Malpighi and Bellaria Hospital, Bologna, Italy, University of Ferrara, Ferrara, Italy*

**Background:** Senile systemic amyloidosis (SSA), due to intramyocardial deposition of wild-type transthyretin (TTR), is often mistaken as hypertensive heart disease (HHD) or hypertrophic cardiomyopathy (HCM). In patients with amyloidotic cardiomyopathy 99mTc-DPD scintigraphy can differentiate between TTR (mutant and wild-type) and primary amyloidosis. We assessed the diagnostic performance of 99mTc-DPD scintigraphy in the non-invasive identification of SSA in a clinical context of elderly patients with unexplained concentric left ventricular (LV) “hypertrophy”.

**Methods:** Patients undergone 99mTc-DPD scintigraphy in 2004-2010 for suspected SSA were included in the analysis. SSA was suspected in patients > 65 years with unexplained concentric LV “hypertrophy” (end-diastolic mean wall thickness > 14 mm) and non-dilated LV, who also presented one or more of: male gender; thickened atrio-ventricular valves; mild pericardial effusion; granular sparkling myocardial appearance; absent or mild LV hypertrophy on ECG. All patients with an already definite diagnosis able to explain LV wall thickness increase were excluded. Myocardial uptake of 99mTc-DPD (740 MBq iv) was semiquantitatively/visually assessed at 5 min/3 h.

**Results:** A total of 62 patients entered the analysis: 32 with SSA, 14 with hereditary TTR-related amyloidosis (ATTR) and exclusively cardiac phenotype and 16 with non-amyloidotic LV “hypertrophy” (9 HCM, 6 HHD, 1 Anderson-Fabry disease). 99mTc-DPD scintigraphy tested positive for myocardial uptake in all cases with TTR amyloidosis (both mutant and wild-type). No myocardial uptake of the tracer was apparent in non-amyloidotic LV hypertrophy. Sensitivity and specificity of positive 99mTc-DPD myocardial tracer uptake (i.e. visual score  $\geq 1$ ) for the diagnosis of SSA and ATTR were both 100%.

**Conclusions:** In elderly patients with unexplained concentric LV “hypertrophy”, 99mTc-DPD scintigraphy can provide a highly accurate technique able to increase the diagnostic performance of echocardiography in the non-invasive identification of TTR-related (both mutant and wild-type) amyloidotic cardiomyopathy.